


# Evidence-Based Medicine



**Editors:** Brian Haynes & Paul Glasziou

## Contents

Purpose and procedure	98	Multicomponent comprehensive care was as effective as usual care for decreasing major bleeding in older patients on warfarin	116
Additional articles abstracted in <i>ACP Journal Club</i>	99	Routine primary immunisation using a longer needle resulted in fewer local reactions in infants	117
<b>EBM notebook</b>		Malaria prophylaxis with atovaquone-proguanil caused fewer gastrointestinal adverse events than chloroquine-proguanil	118
Using patient decision aids to promote evidence-based decision making	100	Psychological therapy reduced depression earlier (4 months) but at 1 year was not better than usual general practitioner care	119
Resource corner	103	Pseudoephedrine and acetaminophen relieved sinus symptoms in the course of the common cold	120
<b>Therapeutics</b>		Ventilation tubes did not improve quality of life in persistent otitis media with effusion	121
Formoterol was more effective than terbutaline when taken as needed for moderate to severe asthma	105	Oedema reduction by mechanical compression improved the healing of foot infection in patients with diabetes mellitus	122
Review: small treatment effect of disodium cromoglycate in childhood asthma is tempered by methodological flaws and publication bias	106	Review: evidence of the effectiveness of discharge planning is equivocal	123
Review: antiplatelet drugs reduce pre-eclampsia, preterm birth, and stillbirth or neonatal death	107	Review: epidural or spinal anaesthesia reduces post-operative mortality and morbidity	124
Review: evidence is sparse and inconclusive for treating and monitoring chronic mild to moderate hypertension in pregnancy	108	<b>Diagnosis</b>	
Combination therapy with candesartan and lisinopril was more effective than monotherapy in type 2 diabetes and hypertension	109	Computed tomography pneumocolon had a low sensitivity for detecting colorectal neoplasms	125
Review: calcium antagonists lead to a higher risk for myocardial infarction, congestive heart failure, and major cardiovascular events	110	<b>Prognosis</b>	
Review: ACE inhibitors, calcium antagonists, and more intensive blood pressure lowering strategies reduce cardiovascular events	111	Subsequent reactions were common and often more serious than the initial reactions of children with peanut allergy	126
Aspirin but not vitamin E prevented cardiovascular events in patients at risk	112	<b>Economics</b>	
Rate and rhythm control showed similar symptom improvement in atrial fibrillation	113	A levonorgestrel releasing intrauterine system was more cost effective than was hysterectomy for menorrhagia	127
Clopidogrel reduced recurrent ischaemic events in patients with previous cardiac surgery more than aspirin	114	<b>Clinical prediction guide</b>	
Vitamin K lowered the international normalised ratio into the therapeutic range in patients receiving warfarin	115	A predictive model using pulmonary function markers identified snorers at low risk for sleep apnoea syndrome	128