Evidence-Based Medicine

Contents

Purpose And Procedure 130	A peer led asthma education programme in adolescents was more effective than no programme for improving quality of
Additional articles obstracted in ACD	life
Additional articles abstracted in ACP Journal Club	Review: some but not all topical antimicrobial agents improve the rate of healing of chronic wounds
EBM notebook	Review: warfarin prevents stroke in non-rheumatic atrial fibrillation but has a higher risk for haemorrhage than other
Creating knowledge management skills in primary care	agents
residents: a description of a new pathway to evidence-based practice in the community	Reduced sodium intake lowered blood pressure and need for antihypertensive medication
Resource corner	Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for
Therapeutics	head lice infestation in children
Review: the levonorgestrel releasing intrauterine system prevents more pregnancies than intrauterine devices (IUDs)	Review: smooth muscle relaxants improve symptoms and reduce pain in irritable bowel syndrome
$\leq 250 \text{ mm}^2 \text{ but not IUDs} > 250 \text{ mm}^2 \dots 137$	A depression management programme reduced depression
Antibiotics did not prevent adverse outcomes after preterm, prelabour rupture of fetal membranes	in frequent users of health care but did not reduce healthcare visits
Antibiotics did not prevent adverse outcomes in women in preterm labour with intact fetal membranes	A depression management programme increased depression free days and costs in depressed frequent users of general health care
Postexposure prophylaxis with inhaled zanamivir was efficacious for household contacts of people with influenza .140	Patient education to encourage graded exercise improved physical functioning in chronic fatigue syndrome156
Oseltamivir reduced new cases of influenza in household contacts of people with clinical or confirmed influenza141	Cognitive behaviour therapy reduced fatigue severity and
Direct observation of tuberculosis treatment did not promote higher cure rates than self administered treatment142	functional impairment in chronic fatigue syndrome157
Review: mefloquine prevents malaria but has adverse effects	Diagnosis
that limit its acceptability in adults not immune to malarial infection	A normal alveolar dead space fraction plus p-dimer assay ruled out suspected pulmonary embolism in the emergency
Incidence of long term backache was similar between	department
women receiving epidural analgesia during labour and those who did not144	The Mini Cog had high sensitivity and specificity for diagnosing dementia in community dwelling older adults159
Radiography led to improved patient satisfaction but increased short term pain in patients with low back pain145	Differential diagnosis
Review: angiotensin converting enzyme inhibitors are beneficial in type 1 diabetes mellitus and microalbuminuria regardless of baseline risk factor status	Differential diagnosis A 2 step procedure determined diagnosis in patients with unexplained weight loss
Review: oral anticoagulants plus antiplatelet agents reduce thromboembolism and all cause mortality for heart valve	
prostheses	Glossary Inside back cover

Now available online at: www.evidence-basedmedicine.com