

Evidence-Based Medicine

Contents

Purpose and procedure	2	Brief psychodynamic interpersonal treatment after deliberate self poisoning reduced suicidal ideation and deliberate self harm	22
Additional articles abstracted in ACP Journal Club	3	Support by another mother and a child-life specialist decreased anxiety in mothers of children with chronic illnesses	23
EBM notebook		Review: cannabinoids and codeine have similar effects on pain relief, but cannabinoids commonly cause psychotropic adverse effects	24
“Double blind, you are the weakest link — goodbye!”	4	Review: cannabinoids control chemotherapy-induced nausea and vomiting but increase the risk of side effects	25
Resource corner	7	Surgery relieved symptoms but decreased survival more than medical treatment in gastro-oesophageal reflux disease	26
Therapeutics		Prognosis	
Review: varicella vaccination is effective in children	9	Progression of disease in HIV infected children slowed after the first year of life	27
Oseltamivir was safe and effective for prophylaxis of influenza in the frail elderly	10	More women with the pelvic girdle syndrome than with other pelvic pain during pregnancy had pelvic pain 2 years after delivery	28
Review: placebo is better than no treatment for subjective continuous outcomes and for treatment of pain	11	Etiology	
Omalizumab reduced inhaled corticosteroid use and exacerbations in childhood allergic asthma	12	Cardiovascular events were increased in patients with high cardiovascular risk who had albuminuria	29
Review: angiotensin-converting enzyme inhibitors reduce the progression of non-diabetic renal disease	13	Quality improvement	
Predischarge coronary angiography was better than exercise testing for reducing hospital use after low-risk chest pain	14	Recall to a general practitioner or to a nurse clinic improved assessment in patients with coronary artery disease	30
Carvedilol reduced mortality and morbidity caused by myocardial infarction in patients with left ventricular dysfunction	15	Clinical prediction guide	
Long term and short term oral anticoagulation treatments were equivalent for venous thromboembolism	16	The Framingham scores overestimated the risk for coronary heart disease in Japanese, Hispanic, and native American cohorts	31
3 months and 1 year of oral anticoagulant treatment were equivalent for idiopathic proximal deep venous thrombosis	17	A clinical prediction model did well in diagnosing paediatric group A β -haemolytic streptococcal pharyngitis	32
Home monitoring service was more effective than usual care in patients with essential hypertension	18	Glossary	Inside back cover
Review: antibiotics modestly increase remission rates in acute bacterial conjunctivitis	19		
Dietary fibre supplementation with psyllium or gum arabic reduced faecal incontinence in community-living adults	20		
17 β -estradiol reduced depressive and somatic symptoms in perimenopausal women	21		

Now available online at:
www.evidence-basedmedicine.com