

Long term and short term oral anticoagulation treatments were equivalent for venous thromboembolism

Pinede L, Ninet J, Duhaut P, et al, for the Investigators of the "Durée Optimale du Traitement AntiVitamines K" (DOTAVK) Study. *Comparison of 3 and 6 months of oral anticoagulant therapy after a first episode of proximal deep vein thrombosis or pulmonary embolism and comparison of 6 and 12 weeks of therapy after isolated calf deep vein thrombosis.* *Circulation* 2001 May 22;103:2453-60.

QUESTION: In patients with proximal deep venous thrombosis (PDVT) or pulmonary embolism, or both, or isolated calf deep venous thrombosis (CDVT), is long term treatment better than short term treatment with oral anticoagulants for preventing recurrences and bleeding complications?

Sources of funding: French Ministry of Health; Hospices Civils de Lyon; Conseil Régional Rhône-Alpes; Association pour la recherche et l'évaluation thérapeutique (APRET).

For correspondence: Dr L Pinede, Department of Internal Medicine, Edouard Herriot Hospital, 69437 Lyons Cedex 03, France. Fax +33 4721 17567.

Long term v short term treatment with oral anticoagulants for venous thromboembolism†

Outcomes at 15 mo	Long term (12 wk)	Short term (6 wk)	RRI (95% CI)	NNH
CDVT				
Recurrence	3%	2%	71% (-65 to 743)	Not significant
Haemorrhage (major, minor, or fatal)	21%	12%	66% (-12 to 216)	Not significant
Major haemorrhage	3%	1%	242% (-50 to 2269)	Not significant
	Long term (6 mo)	Short term (3 mo)		
PDVT/PE				
Recurrence	9%	8%	10% (-37 to 92)	Not significant
Haemorrhage (major, minor, or fatal)	17%	16%	7% (-26 to 57)	Not significant
			RRR (CI)	NNT
Major haemorrhage	2%	3%	60% (-45 to 363)	Not significant

†CDVT = calf deep venous thrombosis; PDVT/PE = proximal deep venous thrombosis or pulmonary embolism, or both. Other abbreviations defined in glossary; RRI, RRR, NNH, NNT, and CI calculated from data in article.

Design

Randomised (allocation concealed*), unblinded,* controlled trial with 15 months of follow up.

Setting

France.

Patients

736 patients (mean age 59 y, 53% women) who had symptomatic PDVT or pulmonary embolism, or both, or symptomatic CDVT confirmed by objective diagnostic tests. Exclusion criteria included pregnancy, breast feeding, vena cava filter implantation, surgical thrombectomy, free-floating thrombus in the inferior vena cava,

active cancer or malignant haematological disease, or a previous venous thromboembolism. 97% of patients completed the study.

Intervention

375 patients (270 with PDVT or pulmonary embolism, or both, and 105 with CDVT) were allocated to short term (3 mo for PDVT-pulmonary embolism, 6 wk for CDVT) oral anticoagulant treatment, and 361 patients (269 with PDVT or pulmonary embolism, or both, and 92 with CDVT) were allocated to long term (6 mo for PDVT-pulmonary embolism, 12 wk for CDVT) treatment with oral anticoagulants. Fluindione was used for oral anticoagulation with dose adjustments to maintain an international normalised ratio of 2.0 to 3.0.

Main outcome measures

Incidence of recurrent venous thromboembolism and haemorrhage (major, minor, or fatal).

Main result

Intention to treat analyses showed that the treatment groups did not differ for incidence rates of recurrent venous thromboembolism and haemorrhage (table).

Conclusions

In patients with proximal deep venous thrombosis or pulmonary embolism, or both, 6 months is equivalent to 3 months of treatment with anticoagulants for preventing recurrences and bleeding complications. Similarly, in patients with calf deep venous thrombosis, 6 weeks is equivalent to 12 weeks of treatment with oral anticoagulants.

*See glossary.

COMMENTARY

5 previous, well-designed, randomised trials have compared different durations of anticoagulation for venous thromboembolism (VTE).¹⁻⁵ 3 of these studies tested whether the duration of anticoagulation could be reduced from 3¹ or 6 months² to only 4 or 6 weeks in patients who, for the most part, had a first episode of VTE and did not have cancer. All 3 studies found that shortening the duration of treatment resulted in about a 2-fold increase in the rate of recurrent VTE after anticoagulants were stopped, without being associated with a convincing reduction in the rate of bleeding. These studies also found that the risk for recurrent VTE was about 4 times higher for "unprovoked" or "idiopathic" VTE than for thrombosis that was provoked by a transient risk factor, such as recent surgery.

The other 2 trials evaluated an extended duration of anticoagulant treatment in 2 groups of patients with VTE who were thought to have a high risk for recurrence: those with a second episode of VTE³ and those with a first unprovoked episode of VTE.⁵ Both studies found that extending oral anticoagulant treatment (international normalised ratio of 2.0 to 3.0) was effective for preventing recurrent VTE; however, long term use was associated with a substantial risk for bleeding. Schulman *et al*³ concluded that after a second episode of thrombosis the risk for recurrent VTE after completing 6 months of treatment (about 6% per year) was not high enough to routinely recommend extended treatment. Our group found that after a first episode of unprovoked thrombosis, the risk for recurrent VTE after completing 3 months of treatment was higher than expected (about 25% during the first year), which justified treatment for longer than 3 months; however, we could not say how much longer than 3 months was required.⁵

continued on next page

3 months and 1 year of oral anticoagulant treatment were equivalent for idiopathic proximal deep venous thrombosis

Agnelli G, Prandoni P, Santamaria MG, et al, and the Warfarin Optimal Duration Italian Trial Investigators. *Three months versus one year of oral anticoagulant therapy for idiopathic deep venous thrombosis*. *N Engl J Med* 2001 Jul 19;345:165–9.

QUESTION: In patients who have had a first episode of idiopathic proximal deep venous thrombosis, what is the long term effectiveness of extending to 1 year the initial 3 month course of oral anticoagulant treatment?

Design

Randomised (unclear allocation concealment*), unblinded,* controlled trial with ≥ 2 years of follow up.

Setting

10 study centres in Italy.

Patients

267 patients (mean age 67 y, 58% men) who had a first episode of symptomatic idiopathic proximal deep venous thrombosis confirmed by compression ultrasonography or venography, and had completed 3 uninterrupted months of oral anticoagulant treatment without a recurrence of thromboembolism or bleeding. Exclusion criteria included anticoagulant treatment for reasons other than venous thromboembolism, major psychiatric disorders, life expectancy < 2 years, and inability to return for follow up. 90% of patients completed the study.

Intervention

After 3 months of treatment with warfarin (97% of patients) or acenocoumarol, 133 patients were allocated to discontinue (3 mo treatment group) and 133 patients to continue (1 y treatment group) oral anticoagulant treatment for 9 additional months. The dose of the oral anticoagulant was adjusted to maintain an international normalised ratio of 2.0 to 3.0.

Main outcome measures

The primary outcome was recurrent deep venous thrombosis. Secondary outcomes were major bleeding and all-cause mortality.

Main results

At ≥ 2 years, intention to treat analyses showed that the groups did not differ for incidence rates of recurrent deep venous thrombosis, major bleeding, and all-cause mortality (table). The study had an 80% power to detect a 50% reduction in the risk of recurrence at the 5% level of significance.

Conclusion

In patients who had a first episode of idiopathic proximal deep venous thrombosis, 3 months was equivalent to 1 year of oral anticoagulant treatment in preventing recurrences at ≥ 2 years.

*See glossary.

Source of funding: not stated.

For correspondence: Professor G Agnelli, Sezione de Medicina Interna e Cardiovascolare, Dipartimento di Medicina Interna, Università di Perugia, Via Enrico dal Pozzo, 06123 Perugia, Italy.

1 year v 3 months' treatment with oral anticoagulants for proximal deep venous thrombosis†

Outcomes at ≥ 2 years	1 year treatment	3 months' treatment	RRR (95% CI)	NNT
Recurrence	16%	16%	1% (–72 to 43)	Not significant
All-cause mortality	5%	5%	1% (–165 to 63)	Not significant
			RRI (CI)	NNH
Major bleeding	3%	2%	99% (–57 to 817)	Not significant

†Abbreviations defined in glossary; RRR, RRI, NNT, NNH, and CI calculated from data in article.

COMMENTARY—continued from previous page

In a mix of patients with provoked and unprovoked VTE, Pinede *et al* found the risk for recurrent VTE to be similar after completing 3 and 6 months of treatment. In patients with a first episode of unprovoked deep venous thrombosis, Agnelli *et al* found a similar risk for recurrent VTE after completing 3 months and 12 months of treatment. In the latter study, this risk was only about 5% per year, much lower than the 25% observed after 3 months of treatment in a similarly defined patient population.⁵

A consistent finding across studies is that 3 months of anticoagulant treatment is adequate for patients with VTE that has been provoked by a major transient risk factor. The subsequent risk for recurrence is low ($< 5\%$ per year); however, uncertainty exists when the risk for recurrence after an unprovoked VTE is as low after 3 months of treatment as it is after 6 months. Although the 2 new studies strongly suggest that 3 months of treatment achieves as low a subsequent risk for recurrent VTE as 6 or 12 months of treatment, neither study had the power to establish this definitively. On the basis (mostly) of the results of our own study,⁵ which is representative of our patient population, I will continue to recommend a minimum of 6 months of anticoagulant treatment for a first unprovoked episode of VTE, provided patients do not have a high risk for bleeding. Current evidence suggests that patients with unprovoked VTE who have been treated for 6 months or longer have a risk for recurrent VTE of about 10% during the year after anticoagulant treatment is stopped. Therefore, once patients have successfully completed 6 months of treatment, I encourage them to remain on long term treatment provided they have a low risk for bleeding and do not find treatment a burden.

Clive Kearon, MB, PhD

Henderson General Hospital, McMaster University, Hamilton, Ontario, Canada

- 1 Research Committee of the British Thoracic Society. Optimum duration of anticoagulation for deep-vein thrombosis and pulmonary embolism. *Lancet* 1992;340:873–6.
- 2 Schulman S, Rhedin AS, Lindmarker P, et al. A comparison of six weeks with six months of oral anticoagulant therapy after a first episode of venous thromboembolism. *N Engl J Med* 1995;332:1661–5.
- 3 Levine MN, Hirsh J, Gent M, et al. Optimal duration of oral anticoagulant therapy: a randomized trial comparing four weeks with three months of warfarin in patients with proximal deep vein thrombosis. *Thromb Haemost* 1995;74:606–11.
- 4 Schulman S, Granqvist S, Holmstrom M, et al. The duration of oral anticoagulant therapy after a second episode of venous thromboembolism. *N Engl J Med* 1997;336:393–8.
- 5 Kearon C, Gent M, Hirsh J, et al. A comparison of three months of anticoagulation with extended anticoagulation for a first episode of idiopathic venous thromboembolism. *N Engl J Med* 1999;340:901–7.