

Evidence-Based Medicine

Editors: Brian Haynes & Paul Glasziou



| | | |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|-----|--|
| Contents | | |
| Purpose and procedure | 98 | |
| Additional articles abstracted in <i>ACP Journal Club</i> | 99 | |
| EBM notebook | | |
| Textbook descriptions of disease — where's the beef? | 100 | |
| Resource review | 103 | |
| Therapeutics | | |
| Review: a home-like birth environment has beneficial effects on labour and delivery | 105 | |
| Review: antibiotics improve maternal and fetal outcomes and are safe in preterm, prelabour rupture of membranes | 106 | |
| Review: interventions that lower cholesterol concentrations or blood pressure in diabetic patients prevent cardiovascular disease | 107 | |
| Review: glycoprotein IIb/IIIa inhibitors reduced death or myocardial infarction in acute coronary syndromes not routinely scheduled for revascularisation | 108 | |
| Treating isolated systolic hypertension prevented major cardiovascular events across strata of risk in older patients | 109 | |
| Review: antiplatelet treatment prevents occlusive vascular events in high risk patients | 110 | |
| Review: aspirin reduces CAD events in people with no history of cardiovascular disease, but it increases gastrointestinal bleeding | 111 | |
| A preoperative smoking intervention decreased postoperative complications in elective knee or hip replacement | 112 | |
| Oral dexamethasone led to fewer treatment failures than did nebulised dexamethasone or placebo in children with mild croup | 113 | |
| Growth rate was greater with fluticasone propionate than with beclomethasone dipropionate in children with chronic asthma | 114 | |
| Cardiac pacing reduced non-accidental falls in older adults with cardioinhibitory carotid sinus hypersensitivity | 115 | |
| Review: intrinsic and environmental risk factor modification reduces falls in elderly people | 116 | |
| Adding intranasal fluticasone to cefuroxime resolved acute rhinosinusitis | 117 | |
| Review: oral and intravaginal agents are equally effective for treatment of uncomplicated vulvovaginal candidiasis | 118 | |
| Verbal advice plus an information leaflet reduced antibiotic use in acute bronchitis | 119 | |
| Low dose budesonide improved asthma control in mild asthma; adding formoterol improved control in corticosteroid treated patients | 120 | |
| Sildenafil improved erectile dysfunction and quality of life in men with comorbid mild to moderate depression | 121 | |
| A multidisciplinary community based rehabilitation programme improved social functioning in severe traumatic brain injury | 122 | |
| Rofecoxib, 25 mg/day, was more effective than rofecoxib, 12.5 mg/day, celecoxib, or acetaminophen in osteoarthritis of the knee | 123 | |
| Review: behavioural interventions plus laxatives are effective for defaecation disorders, but biofeedback does not add benefit | 124 | |
| Naltrexone for 3 or 12 months did not reduce drinking in alcohol dependence | 125 | |
| Diagnosis | | |
| Magnetic resonance direct thrombus imaging was accurate in suspected deep venous thrombosis | 126 | |
| Quality improvement | | |
| A comprehensive annotated reminder tool increased appropriate screening in primary care | 127 | |
| Clinical prediction guide | | |
| A simple scoring system predicted clinical progression in HIV patients receiving highly active antiretroviral therapy | 128 | |