Review: ipratropium is not more effective than β_2 agonists for acute exacerbations of chronic obstructive pulmonary disease

McCrory DC, Brown CD. *Anti-cholinergic bronchodilators versus beta2-sympathomimetic agents for acute exacerbations of chronic obstructive pulmonary disease. Cochrane Database Syst Rev 2002;(4):CD003900.*

QUESTION: In patients with acute exacerbations of chronic obstructive pulmonary disease (COPD), are anticholinergic agents more effective than placebo or β_2 agonists?

Data sources

Studies were identified by searching Medline, EMBASE/Excerpta Medica, CINAHL, and the Cochrane COPD Trials Register, using the terms bronchodilator, ipratropium, and oxitropium; and by scanning bibliographies of relevant studies.

Study selection

Studies were selected if they were randomised controlled trials (RCTs) comparing anticholinergic agents (ipratropium or oxitropium bromide, given by inhalation by nebuliser or metered dose inhaler) with an appropriate control (eg, placebo, other bronchodilating agents, or combination therapies) and assessed adults with a diagnosis of COPD having symptoms of acute exacerbation of COPD. Studies of patients with acute asthma or those receiving ventilation were excluded.

Data extraction

Data were extracted on study quality and methods, participants, interventions, and outcomes (eg, lung function measurements, arterial blood gas measurements, and symptom scores).

Main results

9 RCTs were included. 4 studies compared ipratropium with an inhaled β_2 agonist, and 5 studies compared ipratropium plus a short acting β_2 agonist with a β_2 agonist alone. The most common outcome reported was FEV₁. 4 studies that compared ipratropium bromide with a β_2 agonist showed no difference between groups for change in FEV₁ at 90 minutes (weighted mean difference [WMD] 0.1, 95% CI -0.19 to 0.19). 5 studies that compared ipratropium plus a β_2 agonist with a β_2 agonist alone also showed no difference between groups for change in FEV₁ at 90 minutes (WMD 0.02 l, CI -0.08 to 0.12) and at 24 hours (WMD -0.05 l, CI -0.14 to 0.05). 2 studies that compared ipratropium with a β_2 agonist showed no difference between groups for hypoxia (PaO₂), either in the short or long term.

Conclusion

In patients with acute exacerbations of chronic obstructive pulmonary disease, ipratropium alone or combined

with a short acting β_2 agonist does not increase the degree of bronchodilation more than a short acting β_2 agonist used alone.

COMMENTARY

Current guidelines recommend that an anticholinergic agent be combined with a β_2 adrenergic agonist when inhaled bronchodilator treatment is prescribed for patients with severe or exacerbated COPD. Short acting anticholinergics and short acting β_2 agonists produce similar bronchodilation in patients with stable COPD.¹ Combined treatment produces larger FEV₁ increases than does either agent alone, but the additive effect is modest and the clinical benefit remains uncertain.¹

The meta-analysis by McCrory and Brown shows that anticholinergics and short acting β_2 agonists are similar also when given alone to patients admitted to hospital for COPD exacerbations. However, combined treatment seems to provide no additive bronchodilation during exacerbation, either short (90 mins) or longer term (24 h).

Several reasons are possible for this seeming disparity, although none are readily apparent. Chance can never be fully excluded. Although several hundred patients are included in the meta-analysis, there is a small possibility that FEV₁ differences as large as 100 ml might not have been detected, and changes of this magnitude during exacerbation seem to be clinically meaningful.²

None of the trials included in the meta-analysis attempted to evaluate clinical outcomes. McCrory and Brown suggest the need for a very large trial to better evaluate patient oriented endpoints, but it is doubtful that support will ever be found for such a large, expensive project. Hence, treatment recommendations will continue to be made from the available evidence, which provides little justification for combining an anticholinergic with a β_2 agonist in the treatment of COPD exacerbations. As monotherapy, the 2 drug classes seem to be equally effective, and because both have excellent safety profiles, individual patient preference or cost should dictate the choice.

Dennis E Niewoehner, MD
Veterans Affairs Medical Center
Minneapolis, Minnesota, USA

- 1 In chronic obstructive pulmonary disease, a combination of ipratropium and albuterol is more effective than either agent alone. An 85-day multicenter trial. COMBIVENT Inhalation Aerosol Study Group. *Chest* 1994;**105**:1411-9.
- 2 Niewoehner DE, Collins D, Erbland ML. Relation of FEV₁ to clinical outcomes during exacerbations of chronic obstructive pulmonary disease. Department of Veterans Affairs Cooperative Study Group. *Am J Respir Crit Care Med* 2000;**161**:1201-5.

Source of funding:
Agency for Health Care
Research and Quality.

For correspondence:
Dr D McCrory, Duke
University Medical
Center, Durham, NC,
USA.
douglas.mccrory@
duke.edu

Abstract and
commentary also
appear in ACP Journal
Club.